


Remark on 4.2:


(i) $\vec{F} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$




(ii) $\vec{F} = \begin{pmatrix} y \\ 0 \end{pmatrix}$







(iii) $\vec{F} = \begin{pmatrix} -y \\ x \end{pmatrix}$



$\vec{F}(t) = \begin{pmatrix} r\cos(wt) \\ r\sin(wt) \end{pmatrix}$


$\vec{v} = \begin{pmatrix} -rw\sin \\ rw\cos \end{pmatrix} = w \begin{pmatrix} -y \\ x \end{pmatrix}$

$x = r\cos$

$y = r\sin$

$\rightarrow$ curl den. $(\vec{v}) = 2w$

(vi) $\vec{F} = \frac{1}{x^2+y^2} \begin{pmatrix} -y \\ x \end{pmatrix}$



Stokes thm.: Overview:

Have seen: Green's thm.: (tang. form)


$$\int_C \vec{F} \cdot d\vec{r} = \iint_\Omega \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dA$$



$\vec{F} = \begin{pmatrix} M \\ N \end{pmatrix}$.

Is a thm. in x-y-plane

Goal: Version for Ω a surface in $\mathbb{R}^3$:



$\rightarrow$ Stokes thm. $\int_C \vec{F} \cdot d\vec{r} = \iint_S (\vec{\nabla} \times \vec{F}) \cdot d\vec{S}$

line integral — surface integral of vector field $\vec{\nabla} \times \vec{F}$

$\rightarrow$ need to understand

- surfaces
- surface integrals
- Operator $\vec{\nabla} \times \vec{F}$ (curl)

Surfaces

Def. A parametric representation of a surface is a function:


$$\vec{r} : D \longrightarrow \mathbb{R}^3$$

$$(u,v) \longmapsto \vec{r}(u,v) = \begin{pmatrix} f(u,v) \\ g(u,v) \\ h(u,v) \end{pmatrix},$$

where f, g, h are real valued functions, $D \subset \mathbb{R}^2$.

The set: $S = \{ \vec{r}(u,v) : (u,v) \in D \}$


is called the surface.



Remarks: (i) u, v : parameters

(ii) D : parameter domain


(iii) Relation to curves:



I.e. it is 2-D extension of curve.

Ex: (i) $\vec{r} : \mathbb{R}^2 \longrightarrow \mathbb{R}^3$

$$(u,v) \longmapsto \vec{r}(u,v) = \underbrace{\begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix}}_{=\vec{v}_0} + u \underbrace{\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}}_{\vec{a}} + v \underbrace{\begin{pmatrix} 17 \\ \sqrt{2} \\ -\pi \end{pmatrix}}_{\vec{b}}$$




A plane

(In general: $\vec{r}(u,v) = \vec{v}_0 + u\vec{a} + v\vec{b}$)

(ii) $\vec{r} : [0,1] \times [0, 2\pi] \longrightarrow \mathbb{R}^3$

$$(u,v) \longmapsto \vec{r}(u,v) = \begin{pmatrix} u\cos(v) \\ u\sin(v) \\ u \end{pmatrix}$$




a cone!

Rem.: Usually we use r, φ as parameters: $\vec{r}(r,\varphi) = \begin{pmatrix} r\cos(\varphi) \\ r\sin(\varphi) \\ r \end{pmatrix}$

(iii) **Surfaces of revolution:**

Take a curve in x-z-plane: $\vec{r} : [0,1] \longrightarrow \mathbb{R}^3$


$$u \longmapsto \vec{r}(u) = \begin{pmatrix} u \\ 0 \\ u \end{pmatrix}$$



Then rotate around z-axis: $(v \in [0, 2\pi])$

$$\vec{r}(u,v) = \begin{pmatrix} \cos(v) & -\sin(v) & 0 \\ \sin(v) & \cos(v) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} u \\ 0 \\ u \end{pmatrix} = \begin{pmatrix} u\cos(v) \\ u\sin(v) \\ u \end{pmatrix}$$


again our cone:



(iv) **Sphere:**


Consider the curve in x-z-plane: $\vec{r} : [0, \pi] \longrightarrow \mathbb{R}^3$

$$u \longmapsto \vec{r}(u) = \begin{pmatrix} \sin(u) \\ 0 \\ \cos(u) \end{pmatrix}$$



rotation : $\left(v \in [0, 2\pi] \right)$

$$\vec{r}(u,v) = \begin{pmatrix} \cos(v) & -\sin(v) & 0 \\ \sin(v) & \cos(v) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \sin(u) \\ 0 \\ \cos(u) \end{pmatrix} = \begin{pmatrix} \sin(u)\cos(v) \\ \sin(u)\sin(v) \\ \cos(u) \end{pmatrix}$$



Rem.:

$$\vec{r}(\theta, \varphi) = \begin{pmatrix} \sin(\theta)\cos(\varphi) \\ \sin(\theta)\sin(\varphi) \\ \cos(\theta) \end{pmatrix}$$

θ : polar angle


φ : azimuthal angle

(v) Graphs:

Let $f : \mathbb{R}^2 \longrightarrow \mathbb{R}$


$(x,y) \longmapsto z = f(x,y)$

Then: $\vec{r}(x,y) = \begin{pmatrix} x \\ y \\ f(x,y) \end{pmatrix}$ is a surface : the graph of f.




(everything in one picture !)

Surface Area



$P_0 = \vec{r}(u_0, v_0)$


Patch magnified:



$$\vec{r}_u = \frac{\partial \vec{r}}{\partial u}$$

$$\vec{r}_v = \frac{\partial \vec{r}}{\partial v}$$

Approx. of patch element by parallelogram:



→ Area of patch element:

$$\Delta A \approx |\vec{r}_u \Delta u \times \vec{r}_v \Delta v|$$

$$= |\vec{r}_u \times \vec{r}_v| \, \Delta u \, \Delta v$$

→ Total area $\approx \sum |\vec{r}_u \times \vec{r}_v| \, \Delta u \, \Delta v$

→ **Area:**

$$A = \iint_D |\vec{r}_u \times \vec{r}_v| \, du\, dv$$

Ex: Surface of cone:

$$\vec{r}(u,v) = \begin{pmatrix} u\cos(v) \\ u\sin(v) \\ u \end{pmatrix} \longrightarrow \vec{r}_u = \frac{\partial \vec{r}}{\partial u} = \begin{pmatrix} \cos(v) \\ \sin(v) \\ 1 \end{pmatrix}$$

$$\vec{r}_v = \frac{\partial \vec{r}}{\partial v} = \begin{pmatrix} -u\sin(v) \\ u\cos(v) \\ 0 \end{pmatrix}$$

$$\longrightarrow |\vec{r}_u \times \vec{r}_v| = \left| \begin{pmatrix} \cos(v) \\ \sin(v) \\ 1 \end{pmatrix} \times \begin{pmatrix} -u\sin(v) \\ u\cos(v) \\ 0 \end{pmatrix} \right| = \left| \begin{pmatrix} -u\cos(v) \\ -u\sin(v) \\ u\cos^2(v) + u\sin^2(v) \end{pmatrix} \right|$$

$$= u\sqrt{2}$$

$$\longrightarrow A = \iint_D |\vec{r}_u \times \vec{r}_v| \, du\, dv = \int_{v=0}^{2\pi} \int_{u=0}^{1} \sqrt{2}\, u \, du\, dv = \ldots = \sqrt{2}\pi$$

